



AHA ANTI-AGE LIFT MOISTURISING TREATMENT 6708

Formula

A	<ul style="list-style-type: none">• MONTANOV 68 (Cetearyl alcohol and cetearyl glucoside - SEPPIC)• Isononyl isononanoate• SEPILIFT DPHP (Dipalmitoyl hydroxyproline - SEPPIC)	4.00% 10.00% 0.50%
B	<ul style="list-style-type: none">• Aqua/Water• Potassium lactate (10%)	QS 100% 8.00%
C	<ul style="list-style-type: none">• SEPIGEL 305 (Polyacrylamide and C13-14 isoparaffin and Laureth-7 - SEPPIC)	3.00%
D	<ul style="list-style-type: none">• SEPICIDE HB (Phenoxyethanol/Methylparaben/Ethylparaben /Propylparaben/Butylparaben - SEPPIC)• SEPICIDE CI (Imidazolidinyl urea - SEPPIC)• Parfum/Fragrance• Sodium hydroxide 24%	0.30% 0.20% 0.20% QS pH=5

Method

Melt A at 75°C. Heat B to 75°C. Add A to B, then start the emulsifier (rotor stator). Introduce C. Stop the emulsifier at approximately 60°C. Start cooling and introduce the ingredients of D at around 30°C. Adjust the final pH to 5.

Comments

SEPILIFT DPHP	Liposoluble hydroxyproline carrier of plant origin. With its anti-wrinkle effect, it moisturises and softens the marks of time (efficacy proven in vivo). Firms tissue by stimulating collagen fibre contraction, protects dermal fibres against enzymatic lysis and shows an anti-radical action. Also gives the emulsion a specific feel.
MONTANOV 68	Glucolipid emulsifier of plant origin. Generates liquid crystals in emulsions easily. Reinforces and prolongs the moisturising properties of SEPILIFT DPHP.
SEPIGEL 305	Thickening and emulsifying agent in very easy-to-use liquid presentation (no pre-dispersion or neutralisation). Compatible with AHAs
SEPICIDE HB / CI	Preservative system.



Characteristics

Appearance	white fluid cream.
pH	approx. 5
Viscosity	24,000 mPa.s BROOKFIELD LVT mobile 4 - 6rpm.
Stability	stable at AT/40/50°C stable in thermal cycles (-10°C/+40°C) stable after centrifugation at 50°C

Notes

Fragrance: LIANE (QUEST)
Potassium lactate (10%): PURAC BF/P41 (PURAC)

6708 - SEPPIC – A0001

Since the proposed formulation has not undergone a toxicological study, the handling and use of the proposed products are given as an indication only and in no way bind SEPPIC's responsibility.